Ascension Rx

Lumasiran (Oxlumo & Biosimilar) Injection Order Form

Patient Name: _____ **Date of Birth:** _____

Patient Phone #: _____ **Date:** _____

Therapy Status: ☐ New Start ☐ Continuation ☐ Restart **Last Dose:**_____

Height (inches)	Weight (kg)	Allergies

Diagnosis

ICD-10 Code(s): _____ J Code(s): _____

Labs (Recommended to be done within 30 days of first dose)

- ☐ Labs to be reviewed or ordered by infusion center
  - ☐ Baseline
    - ☐
  - ☐ Maintenance (every _____)
    - ☐
- ☐ Labs reviewed by ordering provider, ok to proceed (Labs last done on _____)
- ☐ Other: _____ ☐

Nursing Order

- ☑ Pregnancy status: If uncertain, obtain HCG urine test for women 11 – 50 years prior to treatment.

Pre-Medications (Not routinely recommended, use only if clinically indicated)

- ☐ Acetaminophen 650 mg PO once
- ☐ Diphenhydramine 25 mg PO once
- ☐ Diphenhydramine 25 mg IV once
- ☐ Diphenhydramine 50 mg PO once
- ☐ Diphenhydramine 50 mg IV once
- ☐ Famotidine 20 mg IV once
- ☐ Famotidine 20 mg PO once
- ☐ Dexamethasone 20 mg IV once
- ☐ Dexamethasone 20 mg PO once
- ☐ Methylprednisolone 125 mg IV once
- ☐ Methylprednisolone 40 mg IV once
- ☐ Other: _____

Ascension Rx

Lumasiran (Oxlumo & Biosimilar) Injection Order Form

Medication Orders

Loading Dose:

☐ Lumasiran 3 mg/kg (_____ mg) SQ once monthly for 3 doses (20 kg and above)

Maintenance Dose:

☐ Lumasiran 3 mg/kg (_____ mg) SQ once every 3 months, beginning 1 month after the last loading dose (20 kg and above)

Refills:
(Refills for one year from date of signature unless indicated)

☐ Zero ☐ For 12 months ☐ _____

To be completed by Infusion Center staff only:

Biosimilar substitution/Product Dispensed: _____ J-Code: _____
Rounded Dose: _____ mg

Rationale for Substitution:

☐ Payer-mandated substitution
☐ Financial stewardship
☐ Patient access / hardship

Pharmacy may substitute a biosimilar based on insurance coverage.
Pharmacy may adjust diluent, volume, and infusion duration in accordance with local practice standards.

Administration Instructions

Administer subcutaneous injection into the abdomen, thigh, or the side or back of the upper arms. Rotate injection sites. Do not inject into scar tissue or areas that are reddened, inflamed, or swollen. If injecting into the abdomen, avoid the area around the navel. If more than one injection is needed for a single dose of lumasiran , the injection sites should be at least 2 cm apart.

Emergency / Hypersensitivity Orders

☑ Notify provider for next steps once patient stabilized
☑ Follow standard orders for management of allergies and anaphylaxis reactions
☑ In absence of standard protocol, follow the orders below:
- ☑ Nursing to stop infusion, maintain airway, place supine, elevate legs, O_2 2 L NC
- ☑ Epinephrine 0.3 mg IM Once PRN anaphylaxis (first-line)
- ☑ Diphenhydramine 25 mg IV q6h PRN itching/hives
- ☑ Hydrocortisone 100 mg IV Once PRN hypersensitivity
- ☑ Methylprednisolone 125 mg IV Once PRN hypersensitivity
- ☑ Famotidine 20 mg IV Once PRN hives/itching
- ☑ Albuterol inhaler 2 puffs Once PRN wheezing/SOB
- ☑ Normal Saline 1000 mL IV bolus Once PRN hypotension

Ascension Rx

Lumasiran (Oxlumo & Biosimilar) Injection Order Form

Infusion Center Staff

Please fax infusion records or discharge summary to: Fax Number _____

Provider Information

Provider Signature: _____ Date: _____

Provider Printed Name: _____ Provider NPI: _____

Provider Address: _____ Phone: _____ Fax: _____

All orders are valid for 1 year from the ordered date unless otherwise indicated