

Mirikizumab-mrkz (OmvoH & Biosimilars) Injection Order Form**Patient Name:** _____**Date of Birth:** _____**Patient Phone #:** _____**Date:** _____**Therapy Status:** ☐ New Start ☐ Continuation ☐ Restart**Last Dose:** _____

Height (inches)	Weight (kg)	Allergies

Diagnosis

ICD-10 Code(s): _____

J Code(s): _____

Labs (Recommended to be done within 30 days of first dose)

- ☐ Labs to be reviewed or ordered by infusion center
- ☒ Baseline
 - ☒ TB screening (QuantIFERON or alternative)
 - ☒ Hepatitis Screening prior to therapy (HBsAg, HepB core Ab total, HCV)
 - ☒ CBC with Differential
 - ☒ CMP
 - ☒ Maintenance (every _q 6 months _____)
 - ☒ CMP
 - ☒ CBC with Differential
- ☐ Labs reviewed by ordering provider, ok to proceed (Labs last done on _____)
- ☐ Other: _____

Provider Information

Live vaccines should not be given concurrently

Nursing Orders

- ☒ Do not administer with an active infection, including clinically important localized infection, or if patient is taking antibiotics for current infection.
- ☒ Pregnancy status: If uncertain, obtain HCG urine test for women 11 – 50 years prior to treatment.

Pre-Medications (Not routinely recommended, use only if clinically indicated)

- | | |
|--|--|
| ☐ Acetaminophen 650 mg PO once | ☐ Dexamethasone 20 mg IV once |
| ☐ Diphenhydramine 25 mg PO once | ☐ Dexamethasone 20 mg PO once |

Mirikizumab-mrkz (Omvo & Biosimilars) Injection Order Form

- | | |
|--|--|
| ☐ Diphenhydramine 25 mg IV once | ☐ Methylprednisolone 125 mg IV once |
| ☐ Diphenhydramine 50 mg PO once | ☐ Methylprednisolone 40 mg IV once |
| ☐ Diphenhydramine 50 mg IV once | ☐ Other: _____ |
| ☐ Famotidine 20 mg IV once | |
| ☐ Famotidine 20 mg PO once | |

IV Site Orders

- ☒ Follow standard protocol for IV site management
- ☒ In absence of standard protocol, follow the orders below:
 - ☒ Sodium chloride 0.9% infusion 250 mL IV at 20 mL/hr PRN (carrier fluid/KVO)
 - ☒ Start peripheral IV or access central line for infusion
 - ☒ Sodium chloride 0.9% 10 mL flush to IV site as needed
 - ☒ Heparin 50 units/5 mL flush to IV site as needed (central line only)
 - ☒ Heparin 500 units/5 mL flush to IV site as needed (central line only)
 - ☒ May use alteplase 2 mg intracatheter as needed for negative blood return of IV site

Medication Orders

Induction Dose ☐ Mirikizumab-mrkz 300 mg IV over 30 minutes once on week 0, 4, 8 (Ulcerative Colitis) ☐ Mirikizumab-mrkz 900 mg IV over 90 minutes once on week 0, 4, 8 (Crohn's Disease)
To be completed by Infusion Center staff only: Biosimilar substitution/Product Dispensed: _____ J-Code: _____ Rounded Dose: _____ mg
Rationale for Substitution: ☐ Payer-mandated substitution ☐ Financial stewardship ☐ Patient access / hardship
Pharmacy may substitute a biosimilar based on insurance coverage. Pharmacy may adjust diluent, volume, and infusion duration in accordance with local practice standards.

Administration Instructions

Monitor patient for 30 minutes after the first infusion. Monitoring not needed for subsequent infusions unless a reaction has occurred.

Emergency / Hypersensitivity Orders

- ☒ Notify provider for next steps once patient stabilized
- ☒ Follow standard orders for management of allergies and anaphylaxis reactions
- ☒ In absence of standard protocol, follow the orders below:
 - ☒ Nursing to stop infusion, maintain airway, place supine, elevate legs, O₂ 2 L NC
 - ☒ Epinephrine 0.3 mg IM Once PRN anaphylaxis (first-line)

Mirikizumab-mrkz (Omvo® & Biosimilars) Injection Order Form

- ☒ Diphenhydramine 25 mg IV q6h PRN itching/hives
- ☒ Hydrocortisone 100 mg IV Once PRN hypersensitivity
- ☒ Methylprednisolone 125 mg IV Once PRN hypersensitivity
- ☒ Famotidine 20 mg IV Once PRN hives/itching
- ☒ Albuterol inhaler 2 puffs Once PRN wheezing/SOB
- ☒ Normal Saline 1000 mL IV bolus Once PRN hypotension

Infusion Center Staff

Please fax infusion records or discharge summary to: Fax Number _____

Provider Information

Provider Signature: _____ Date: _____

Provider Printed Name: _____ Provider NPI: _____

Provider Address: _____ Phone: _____ Fax: _____

All orders are valid for 1 year from the ordered date unless otherwise indicated