Ascension Rx

Pegloticase (Krystexxa & Biosimilars) Injection Order Form

Patient Name: _____ **Date of Birth:** _____

Patient Phone #: _____ **Date:** _____

Therapy Status: ☐ New Start ☐ Continuation ☐ Restart **Last Dose:**_____

Height (inches)	Weight (kg)	Allergies

Diagnosis

ICD-10 Code(s): _____ J Code(s): _____

Labs (Recommended to be done within 30 days of first dose)

- ☐ Labs to be reviewed or ordered by infusion center
  - ☑ Baseline
    - ☐ CMP
    - ☐ CBC with Differential
    - ☑ Uric Acid (24-48 hours prior each infusion)
    - ☐ BNP
    - ☑ G6PD
  - ☑ Maintenance (every 2 weeks)
    - ☐ CMP
    - ☐ CBC with Differential
    - ☑ Uric Acid (24-48 hours prior to each infusion)
    - ☐ BNP
- ☐ Labs reviewed by ordering provider, ok to proceed (Labs last done on _____)
- ☐ Other: _____

Provider Information

All oral antihyperuricemic agents should be discontinued prior to initiating pegloticase and should not be initiated during the course of therapy

Gout flare prophylaxis with NSAIDS or colchicine should be initiated 1 week prior to pegloticase and continued for the first 6 months of therapy

Nursing Order

- ☑ Do not administer if Uric Acid levels increase to above 6 mg/dL, particularly when 2 consecutive levels above 6 mg/dL are observed.

Ascension Rx

Pegloticase (Krystexxa & Biosimilars) Injection Order Form

☑ Pregnancy status: If uncertain, obtain HCG urine test for women 11 – 50 years prior to treatment.

Pre-Medications

☐ Acetaminophen 650 mg PO once

☑ Diphenhydramine 25 mg PO once

☐ Diphenhydramine 25 mg IV once

☐ Diphenhydramine 50 mg PO once

☐ Diphenhydramine 50 mg IV once

☐ Famotidine 20 mg IV once

☐ Famotidine 20 mg PO once

☐ Dexamethasone 20 mg IV once

☐ Dexamethasone 20 mg PO once

☑ Methylprednisolone 125 mg IV once

☐ Methylprednisolone 40 mg IV once

☐ Other: _____

IV Site Orders

☑ Follow standard protocol for IV site management

☑ In absence of standard protocol, follow the orders below:

- ☑ Sodium chloride 0.9% infusion 250 mL IV at 20 mL/hr PRN (carrier fluid/KVO)
- ☑ Start peripheral IV or access central line for infusion
- ☑ Sodium chloride 0.9% 10 mL flush to IV site as needed
- ☑ Heparin 50 units/5 mL flush to IV site as needed (central line only)
- ☑ Heparin 500 units/5 mL flush to IV site as needed (central line only)
- ☑ May use alteplase 2 mg intracatheter as needed for negative blood return of IV site

Medication Orders

☑ Pegloticase 8 mg IV once over a period of no less than 120 minutes every 2 weeks

To be completed by Infusion Center staff only:

Biosimilar substitution/Product Dispensed: _____ J-Code: _____
Rounded Dose: _____ mg

Rationale for Substitution:

☐ Payer-mandated substitution

☐ Financial stewardship

☐ Patient access / hardship

Pharmacy may substitute a biosimilar based on insurance coverage.
Pharmacy may adjust diluent, volume, and infusion duration in accordance with local practice standards.

Administration Instructions

Observe for at least 1 hour after completion of each infusion.

Emergency / Hypersensitivity Orders

☑ Notify provider for next steps once patient stabilized

09/18/2026

Ascension Rx

Pegloticase (Krystexxa & Biosimilars) Injection Order Form

- [x] Follow standard orders for management of allergies and anaphylaxis reactions
- [x] In absence of standard protocol, follow the orders below:
  - [x] Nursing to stop infusion, maintain airway, place supine, elevate legs, O_2 2 L NC
  - [x] Epinephrine 0.3 mg IM Once PRN anaphylaxis (first-line)
  - [x] Diphenhydramine 25 mg IV q6h PRN itching/hives
  - [x] Hydrocortisone 100 mg IV Once PRN hypersensitivity
  - [x] Methylprednisolone 125 mg IV Once PRN hypersensitivity
  - [x] Famotidine 20 mg IV Once PRN hives/itching
  - [x] Albuterol inhaler 2 puffs Once PRN wheezing/SOB
  - [x] Normal Saline 1000 mL IV bolus Once PRN hypotension

Infusion Center Staff

Please fax infusion records or discharge summary to: Fax Number _____

Provider Information

Provider Signature: _____ Date: _____

Provider Printed Name: _____ Provider NPI: _____

Provider Address: _____ Phone: _____ Fax: _____

All orders are valid for 1 year from the ordered date unless otherwise indicated