

Ublituximab-xiyy (Briumvi & Biosimilars) Injection Order Form**Patient Name:** _____**Date of Birth:** _____**Patient Phone #:** _____**Date:** _____**Therapy Status:** ☐ New Start ☐ Continuation ☐ Restart**Last Dose:** _____

Height (inches)	Weight (kg)	Allergies

Diagnosis

ICD-10 Code(s): _____

J Code(s): _____

Labs (Recommended to be done within 30 days of first dose)

- ☐ Labs to be reviewed or ordered by infusion center
- ☐ Baseline
 - ☒ CMP
 - ☐ CBC with Differential
 - ☐ Quantitative Serum Immunoglobulins (IgG, IgM, IgA)
 - ☒ Hepatitis Screening prior to therapy (HBsAg, HepB core Ab total, HCV)
 - ☐ Maintenance (every _____)
 - ☐ CMP
 - ☐ CBC with Differential
- ☐ Labs reviewed by ordering provider, ok to proceed (Labs last done on _____)
- ☐ Other: _____

Provider Information

Live vaccines should not be given concurrently

Nursing Order

- ☒ Do not administer Brriumvi with an active infection (including active hepatitis B virus), including clinically important localized infection.
- ☒ Pregnancy status: If uncertain, obtain HCG urine test for women 11 – 50 years prior to treatment.

Pre-Medications

- | | |
|---|---|
| ☐ Acetaminophen 650 mg PO once | ☐ Dexamethasone 20 mg IV once |
| ☒ Diphenhydramine 25 mg PO once | ☐ Dexamethasone 20 mg PO once |
| ☐ Diphenhydramine 25 mg IV once | ☒ Methylprednisolone 125 mg IV once |

Ublituximab-xiyy (Briumvi & Biosimilars) Injection Order Form

- | | |
|--|---|
| ☐ Diphenhydramine 50 mg PO once | ☐ Methylprednisolone 40 mg IV once |
| ☐ Diphenhydramine 50 mg IV once | ☐ Other: _____ |
| ☐ Famotidine 20 mg IV once | |
| ☐ Famotidine 20 mg PO once | |

IV Site Orders

- ☒ Follow standard protocol for IV site management
- ☒ In absence of standard protocol, follow the orders below:
 - ☒ Sodium chloride 0.9% infusion 250 mL IV at 20 mL/hr PRN (carrier fluid/KVO)
 - ☒ Start peripheral IV or access central line for infusion
 - ☒ Sodium chloride 0.9% 10 mL flush to IV site as needed
 - ☒ Heparin 50 units/5 mL flush to IV site as needed (central line only)
 - ☒ Heparin 500 units/5 mL flush to IV site as needed (central line only)
 - ☒ May use alteplase 2 mg intracatheter as needed for negative blood return of IV site

Medication Orders

Initial Dose

- ☐ Ublituximab 150 mg IV once over 4 hours followed by 450 mg IV once over 1 hour on day 15

Maintenance Infusions:

- ☐ Ublituximab 450 mg IV once every 24 weeks x 2 doses

To be completed by Infusion Center staff only:

Biosimilar substitution/Product Dispensed: _____ J-Code: _____
Rounded Dose: _____ mg

Rationale for Substitution:

- ☐ Payer-mandated substitution
- ☐ Financial stewardship
- ☐ Patient access / hardship

Pharmacy may substitute a biosimilar based on insurance coverage.

Pharmacy may adjust diluent, volume, and infusion duration in accordance with local practice standards.

Administration Instructions

Observe for at least 1 hour after completion of Infusions #1 and #2. Post-infusion monitoring for subsequent infusions and beyond is at physician discretion, unless a reaction or hypersensitivity has previously occurred with any prior infusion.

- **First Infusion (150 mg):** Begin infusion at 10 mL/hr for 30 minutes, then increase to 20 mL/hr for 30 minutes, then increase to 35 mL/hr for 60 minutes, then increase to 100 mL/hr for the remaining 2 hours
- **Second Infusion (450 mg):** Begin infusion at 100 mL/hr for the first 30 minutes, then increase to 400 mL/hr for the remaining 30 minutes
- **Subsequent Infusions (450 mg):** Begin infusion at 100 mL/hr for the first 30 minutes, then increase to 400 mL/hr for the remaining 30 minutes

Ublituximab-xiiy (Briumvi & Biosimilars) Injection Order Form

Emergency / Hypersensitivity Orders

- ☒ Notify provider for next steps once patient stabilized
- ☒ Follow standard orders for management of allergies and anaphylaxis reactions
- ☒ In absence of standard protocol, follow the orders below:
 - ☒ Nursing to stop infusion, maintain airway, place supine, elevate legs, O₂ 2 L NC
 - ☒ Epinephrine 0.3 mg IM Once PRN anaphylaxis (first-line)
 - ☒ Diphenhydramine 25 mg IV q6h PRN itching/hives
 - ☒ Hydrocortisone 100 mg IV Once PRN hypersensitivity
 - ☒ Methylprednisolone 125 mg IV Once PRN hypersensitivity
 - ☒ Famotidine 20 mg IV Once PRN hives/itching
 - ☒ Albuterol inhaler 2 puffs Once PRN wheezing/SOB
 - ☒ Normal Saline 1000 mL IV bolus Once PRN hypotension

Infusion Center Staff

Please fax infusion records or discharge summary to: Fax Number _____

Provider Information

Provider Signature: _____ Date: _____

Provider Printed Name: _____ Provider NPI: _____

Provider Address: _____ Phone: _____ Fax: _____

All orders are valid for 1 year from the ordered date unless otherwise indicated